

CHECKLIST

New data regarding the distribution and status of the herpetofauna from urban and peri-urban habitats in the city of Pitești, Argeș county (Romania)

Ionuț-Cătălin Petreanu¹

¹ Faculty of Biology, Alexandru Ioan Cuza University of Iași, Blvd. Carol I nr. 20A, 700505, Iași, Romania

Corresponding author: Ionuț-Cătălin Petreanu (icpetreanu@yahoo.com)

Received 3 August 2023 | Accepted 20 November 2023 | Published 31 December 2023

Citation: Petreanu I-C (2023) New data regarding the distribution and status of the herpetofauna from urban and peri-urban habitats in the city of Pitești, Argeș county (Romania). Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa" 66(2): 307–335. <https://doi.org/10.3897/travaux.66.e110472>

Abstract

Identifying urban and peri-urban areas inhabited by species of reptiles and amphibians represents the first step in understanding how herpetofauna can adapt to anthropogenic factors. This is particularly true for regions where sampling biases have left significant gaps in our knowledge of fauna distribution, such as Romania. The aim of the study is to show my preliminary observations on the herpetofauna and its use of habitats in the urban and peri-urban environments from the city of Pitești, Argeș County, located in southern Romania. I identified nine species of amphibians (*Salamandra salamandra*, *Lissotriton vulgaris*, *Triturus cristatus*, *Bombina variegata*, *Bufo bufo*, *Bufotes viridis*, *Hyla orientalis*, *Pelophylax ridibundus* and *Rana dalmatina*) and nine species of reptiles (*Emys orbicularis*, *Trachemys scripta*, *Lacerta agilis*, *Lacerta viridis*, *Podarcis muralis*, *Anguis colchica*, *Coronella austriaca*, *Natrix natrix* and *Natrix tessellata*). The most abundant species in urban environments were *Podarcis muralis* and *Bufotes viridis*, with other species naturally present in the same habitats being *Lacerta viridis*, *Hyla orientalis* and *Bufo bufo*. In an artificial urban pond there were present *Emys orbicularis* and the invasive species *Trachemys scripta* (together with other exotic turtles belonging to other genera) as a result of translocation and releases. When taking into account the peri-urban areas as well, the most abundant species in the study area were *Bombina variegata*, *Rana dalmatina* and *Bufotes viridis*, the first two being limited to the woodland area. The most widespread species were *Hyla orientalis*, *Bufotes viridis*, *Lacerta viridis*, *Podarcis muralis* and *Natrix natrix*. These widespread species were present throughout all the study area. I have also recorded some species of reptiles with very few observations in the Argeș County: *Natrix tessellata* and *Trachemys scripta*. Finally, I identified anthropogenic factors that negatively impact the herpetofauna, such as habitat loss, direct persecution, road mortality, invasive species and poaching.

Keywords

herpetofauna, urban and peri-urban environments, urbanization, reptiles, amphibians, Argeş.

Introduction

Reptiles and amphibians comprise the two vertebrate groups currently most vulnerable and under global population decline (Gibbons et al. 2000; Stuart et al. 2004). These groups exhibit many intrinsic characteristics that increase their proneness to extinction, and lots of anthropogenic threats affect them, such as habitat loss (Mayani-Parás et al. 2019), climate change (Bickford et al. 2010), invasive species, new diseases spread by humans, poaching and direct persecution as a result of public fear, myths or misbeliefs (Fuhn 1969; Marshall et al. 2020). Furthermore, many species of wildlife that live in urban environments and densely populated areas, so it is important to document them and ensure the best preservation of the biodiversity possible, urbanization being part of the habitat loss phenomenon (Hamer and McDonnell 2010; Farinha-Marques et al. 2011). Although urbanization contributes greatly to habitat loss, some species of reptiles and amphibians, including some threatened and protected, are known to occur within urbanized environments (Delaney 2021). Therefore, it is important to explore how the herpetofauna is utilizing the urban environments and habitats, how certain anthropic activities affect it and what measures of urban habitat management should be taken to ensure the preservation of these populations. The Argeş county, situated in Southern Romania is very poor in data regarding the species of reptiles and amphibians, due to the lack of studies conducted, most of the articles on the herpetofauna being focused on the area of the montane regions in the North of the county (Fuhn 1960; Fuhn and Vancea 1961; Strugariu et al. 2009; Covaciu-Marcov et al. 2010, 2014; Iftime and Iftime 2014), but none in the central area where Piteşti is located. This study brings a needed series of data missing from our knowledge on the Romanian fauna, as well as shedding more light on the effects urbanization and expansion of human-inhabited areas have on the local populations of reptiles and amphibians. This work represents the first inventory and mapping survey of the herpetofauna of the urban and peri-urban environments (with some notes regarding natural habitats) of the city of Piteşti (Argeş county), Southern Romania. My study represents a necessary first step for future management and conservation efforts of the area, and a general contribution to the mapping of the Romanian herpetofauna, where sampling biases have left major gaps in knowledge and numerous uncharted areas (Cogălniceanu et al. 2013).

Material and methods

The survey was carried out in the city of Piteşti and its surrounding areas. Piteşti is a city in the center of Argeş County, being surrounded by the Trivale Forest and the

Argeș River, including part of the Natura 2000 site ROSPA0062 (Figure 1). The altitude varies between 250 m and 400 m. I conducted field surveys between 2018 and 2021, mostly by making repeated visits to certain locations that presented favourable habitats for reptiles and amphibians (Edgar et al. 2010; Török et al. 2013) and using visual and auditory transects and active searches for animals (Török et al. 2013). However, I also considered old records, occasional observations from locals and biologists, as well as data from the collections of the Argeș County Museum and the University of Pitești. I visually identified animals without capture as often as possible. Only such specimens that needed rescue/rehabilitation were captured - see text below. I captured animals either by hand or using a net. To identify turtles basking at great distances, I used a pair of 10x42 Opticron binoculars and a Nikon D5600 camera with a 70–300 mm lens. The habitats were varied: forest, river, lake, parks, urban areas etc., and so, the searching methods were adapted accordingly. I visited the rest of the habitats mostly during spring and summer months early in the day, especially during the 07:30–12:00 interval, and looked for reptiles basking and picked up logs, stones, vegetation and garbage. I also identified different amphibians by their call, mostly at night in spring. The identifications were made with the aid of Speybroeck et al. 2016 and Cogălniceanu 2002.




Figure 1. The study area (source: ArcGIS 10.4 basemaps).

Results and discussion

During my research I identified nine species of amphibians: *Salamandra salamandra*, *Lissotriton vulgaris*, *Triturus cristatus*, *Bombina variegata*, *Bufo bufo*, *Bufotes viridis*, *Hyla orientalis*, *Pelophylax ridibundus* and *Rana dalmatina*, belonging to five families (Salamandridae, Bombinatoridae, Bufonidae, Hylidae and Ranidae) and nine species of reptiles: *Emys orbicularis*, *Trachemys scripta*, *Lacerta agilis*, *Lacerta viridis*, *Podarcis muralis*, *Anguis colchica*, *Coronella austriaca*, *Natrix natrix* and *Natrix tessellata*, belonging to five families (Emyidae, Lacertidae, Anguidae, Colubridae and Natricidae).

Peak activity was observed in September with 13 species recorded, followed May and August, with 12 species each. The monthly breakdown of the species observed is shown in the table below (Table 1). One curious and late observation was that of a *Natrix natrix* in early December 2019 (Fuciu Cătălin, pers. communication) (Table 1).

The historical records from the study area were gathered from the specimen collection of the University of Pitești and the Argeș County Museum. It is to be noted that the exact location is almost never given, and instead the label only mentions a locality or a vague, general area. For this reason, the historical observations are not represented on the maps (those only containing the sightings within the study period), instead choosing to put into a table as follows in Table 2.

Out of all the species observed, seven were found in both urban and peri-urban habitats (*Bufo bufo*, *Bufotes viridis*, *Hyla orientalis*, *Lacerta viridis*, *Podarcis muralis* and *Emys orbicularis* and the invasive species *Trachemys scripta*, the last two being present in urban environments as a result of translocation, artificial introduction and human abandonment (park administration, pers. communication)) and the rest exclusively in peri-urban habitats (Table 3). When it comes to urban areas with very few natural elements, the most abundant species were *Podarcis muralis* and *Bufotes viridis*, both found hiding in crevices in concrete and coming out to sun, for hunting, as well as mating.

The most abundant amphibian species were *Bombina variegata*, *Rana dalmatina* and *Bufotes viridis*, while the most abundant reptile species was *Podarcis muralis*. Despite being the most abundant, these species of amphibians were not the most widespread, *Bombina variegata* and *Rana dalmatina* being restricted to the woodland area (Figure 2). The most widespread amphibian species according to the distribution maps created are *Bufotes viridis* and *Hyla orientalis*, while the most widespread reptile species are *Podarcis muralis*, *Lacerta viridis* and *Natrix natrix*. These widespread species were remarkable in being present through the studied area in diverse habitats, be it even insular recordings.

Table 1. Observed presence of each species in the study area by each month.

	January	February	March	April	May	June	July	August	September	October	November	December
<i>Salamandra salamandra</i>			+				+		+(U)	+		
<i>Triturus cristatus</i>		+(M)	+(M)	+	+						+(M)	
<i>Bombina variegata</i>				+	+		+	+	+	+		
<i>Bufo bufo</i>					+		+	+	+			
<i>Bufo viridis</i>						+	+	+	+	+	+	
<i>Hyla orientalis</i>				+(M)		+(Băjan Eliana, pers. communication)	+	+				
<i>Pelophylax sp.</i>				+	+	+	+	+	+			
<i>Rana dalmatina</i>			+	+	+		+	+	+	+	+(M)	
<i>Enys orbicularis</i>			+(D)		+	+	+		+(M)			
<i>Trachemys scripta</i>						+			+			
<i>Lacerta agilis</i>			+			+		+	+(M)			
<i>Lacerta viridis</i>				+	+	+	+	+	+			
<i>Podarcis muralis</i>		+	+	+	+		+	+	+	+	+	
<i>Anguis colchica</i>					+	+		+	+			
<i>Coronella austriaca</i>					+			+	+(M)			
<i>Natrix natrix</i>			+		+	+(M)	+	+				+(Fuciu Cătălin, pers. communication)
<i>Natrix tessellata</i>										+(D)		

“+” – at least one individual of the respective species was observed at least once in the study area during the said month in the course of the study
“+(M)” - the species was not observed by the author during the said month, but there are historical specimens collected in the study area from the said month in the collection of the Argeș County Museum
“+(U)” - the species was not observed by the author during the said month, but there are historical specimens collected in the study area from the said month in the collection of the University of Pitești
“+(D)” - the species was not observed by the author personally in the study area during the said month, but the sightings were provided from public uploads on various databases and social media platforms

Table 2. The inventory of reptile and amphibian specimens in the collections of the Argeş County Museum and the University of Piteşti

Institution	Species	Number of individuals	Date	Location	Collector	Notes
Argeş County Museum	<i>Natrix natrix</i>	1	June 1965	Piteşti AG	Mihăilescu Gh.	adult
Argeş County Museum	<i>Coronella austriaca</i>	1	September 15 1969	Bascov AG	Marius Gh.	adult
Argeş County Museum	<i>Lacerta viridis</i>	1	June 26 1972	Trivale Forest, Piteşti AG	Fusea N.	adult male
Argeş County Museum	<i>Natrix natrix</i>	1	1972	Piteşti AG	?	adult
Argeş County Museum	<i>Anguis colchica</i>	1	September 24 1978	Trivale Forest, Piteşti AG	Mătieş M.	adult; 380 mm
Argeş County Museum	<i>Lacerta viridis</i>	1	July 1977	Trivale Forest	Fusea N.	juvenile
Argeş County Museum	<i>Emys orbicularis</i>	2	September 1976	Ştefăneşti Zăvoi	Mătieş M.	adult female; adult male
Argeş County Museum	<i>Triturus cristatus</i>	1	March 20 1972	Trivale Forest, Piteşti AG	Fusea N.	adult
Argeş County Museum	<i>Triturus cristatus</i>	8	November 7 1972	Trivale Forest, Piteşti AG	Fusea N.	6 adult males; 2 adult females
Argeş County Museum	<i>Rana dalmatina</i>	2	November 7 1972	Trivale Forest, Piteşti AG	Fusea N.	adult male; adult female
Argeş County Museum	<i>Triturus cristatus</i>	12	February 20 1973	Trivale Forest, Piteşti AG	Fusea N.	8 adult males; 4 adult females
Argeş County Museum	<i>Hyla orientalis</i>	3	April 19 1973	Trivale Forest, Piteşti AG	Fusea N.	adults
Argeş County Museum	<i>Rana dalmatina</i>	2	April 19 1973	Trivale Forest	Fusea N.	adult male; adult female
Argeş County Museum	<i>Bombina variegata</i>	6	April 19 1973	Trivale Forest	Fusea N.	2 females; 4 unsexed
Argeş County Museum	<i>Triturus cristatus</i>	5	April 19 1973	Trivale Forest	Fusea N.	females
Argeş County Museum	<i>Pelophylax ridibundus</i>	1	July 10 1973	Trivale Forest	Fusea N.	female

Table 2. (continued)

Institution	Species	Number of individuals	Date	Location	Collector	Notes
Argeș County Museum	<i>Pelophylax sp.</i>	?	April 4 1970	Pitești AG	Mătiș M.	spawn
Argeș County Museum	<i>Pelophylax ridibundus</i>	7	September 20 1974	Bascov Lake	Mătiș M.	unsexed adults
Argeș County Museum	<i>Pelophylax ridibundus</i>	9	September 21 1974	Lunca Golești	Mătiș M.	unsexed adults
Argeș County Museum	<i>Rana dalmatina</i>	1	September 20 1974	Bascov Lake	Mătiș M.	juvenile
Argeș County Museum	<i>Rana dalmatina</i>	2	1977	Trivale Forest	?	juveniles
Argeș County Museum	<i>Triturus cristatus</i>	2	April 25 1974	Pitești AG	Fusea N.	unsexed
Argeș County Museum	<i>Lacerta viridis</i>	1	July 1977	Trivale Forest	Fusea N.	unsexed
Argeș County Museum	<i>Pelophylax ridibundus</i>	1	August 1969	Pitești AG	Balmuș I.	unsexed
Argeș County Museum	<i>Pelophylax ridibundus</i>	1	May 5 2018	Golești AG	Enescu L.	?
University of Pitești	<i>Triturus cristatus</i>	2	May 1968	Pitești AG	Stănescu Gh.	male and female
University of Pitești	<i>Triturus cristatus</i>	1	May 1968	Pitești AG	Stănescu Gh.	female
University of Pitești	<i>Pelophylax esculentus</i>	2	May 1965	Pitești AG	Stănescu Gh.	unsexed
University of Pitești	<i>Rana dalmatina</i>	1	October 1969	Trivale Forest	Stănescu Gh.	unsexed
University of Pitești	<i>Bufoles viridis</i>	1	September 1964	Pitești AG	Stănescu Gh.	unsexed
University of Pitești	<i>Hyla orientalis</i>	1	June 1966	Trivale Forest	Stănescu Gh.	unsexed
University of Pitești	<i>Bufoles viridis</i>	1	1964	Pitești AG	Stănescu Gh.	unsexed
University of Pitești	<i>Salamandra salamandra</i>	2	September 1968	Trivale Forest	Stănescu Gh.	unsexed
University of Pitești	<i>Bombina variegata</i>	1	October 1970	Pitești AG	Stănescu Gh.	unsexed
University of Pitești	<i>Lacerta agilis</i>	1	September 1968	Pitești AG	Stănescu Gh.	Unsexed

Table 3. Use of urban and peri-urban habitats by each species

Species	Habitat
<i>Salamandra salamandra</i>	Peri-urban habitats
<i>Lissotriton vulgaris</i>	Peri-urban habitats
<i>Triturus cristatus</i>	Peri-urban habitats
<i>Bombina variegata</i>	Peri-urban habitats
<i>Bufo bufo</i>	Peri-urban and urban habitats
<i>Bufotes viridis</i>	Peri-urban and urban habitats
<i>Hyla orientalis</i>	Peri-urban and urban habitats
<i>Pelophylax sp.</i>	Peri-urban habitats
<i>Rana dalmatina</i>	Peri-urban habitats
<i>Emys orbicularis</i>	Peri-urban and urban habitats
<i>Trachemys scripta</i>	Peri-urban and urban habitats
<i>Lacerta agilis</i>	Peri-urban habitats
<i>Lacerta viridis</i>	Peri-urban and urban habitats
<i>Podarcis muralis</i>	Peri-urban and urban habitats
<i>Anguis colchica</i>	Peri-urban habitats
<i>Coronella austriaca</i>	Peri-urban habitats
<i>Natrix natrix</i>	Peri-urban habitats
<i>Natrix tessellata</i>	Peri-urban habitats




Figure 2. Observed species of reptiles and amphibians by number of record points. It should be noted that this graphic only shows the points where the species was recorded. The abundance is not always accurately represented as in some record points there have been observed dozens of individuals at a time.

Checklist

Class Amphibia

Order Urodela

Family Salamandridae

Salamandra salamandra (Linnaeus, 1758)

The fire salamander was identified as larvae in 2 locations in the Trivale Forest, and an adult photographed by a local in his garden at the edge of the said forest. The larvae were found in small ponds on the forest floor in early July 2020 and early August 2020. Another adult was recorded on October 24th 2021 dead on a trail in the same forest with the cause of death unknown, but seemingly crushed. The habitat these animals were found in is a deciduous forest, at an altitude of 320 m and 400 m. (Figure 3)

Lissotritton vulgaris (Linnaeus, 1758)

The smooth newt was found in three locations: two separate individuals in ponds at the edge of the forest with very thick vegetation as adults in the aquatic phase, one of them being a breeding male. The animals were both observed in April 2021. One more larva was observed on May 23rd 2020 in a stream. (Figure 4)




Figure 3. Distribution map of *Salamandra salamandra* in the studied area.




Figure 4. Distribution map of *Lissotriton vulgaris* in the studied area.

Triturus cristatus (Laurenti, 1768)

The great crested newt was found in only two locations. One adult female in the aquatic phase was found in April 2021 in a stream flowing at the edge of the Trivale Forest and another adult was found in early April 2018 dead on the road, run over on the street going through the forest next to a pond where the smooth newt was also present. (Figure 5)

Order Anura

Family Bombinatoridae

Bombina variegata (Linnaeus, 1758)

The yellow-bellied toad was found in many locations in the Trivale Forest in small, usually temporary, ponds and streams. It is an abundant species, very abundant in most temporary water bodies in the woods. Its range was, unsurprisingly, restricted to the forest. All life stages were observed. (Figure 6)




Figure 5. Distribution map of *Triturus cristatus* in the studied area.




Figure 6. Distribution map of *Bombina variegata* in the studied area.

Family Bufonidae
Bufo bufo (Linnaeus, 1758)

The common toad was mostly recorded in and around the forest, with a few observations in parts of the urban habitat neighbouring the woods, taking shelter in cracks in the concrete. (Figure 7)

Bufo viridis (Laurenti, 1768)

Although the green toad was also found at the edge of the forest, this amphibian was observed many times in urban and suburban areas, using cracks in the pavement as shelter and coming out to feed at night. A lot of the specimens found were roadkills during the migration towards the breeding spots. (Figure 8)

Family Hylidae
Hyla orientalis Bedriaga, 1890

I managed to find the eastern tree frog in a variety of habitats, the most common one being the forest edge, but it was also recorded in suburban and urban areas and in the reed beds on the banks of the Prundu Lake, on the Argeş River. In July and August 2018 I recorded an interesting breeding site of this species: a pair (or more)




Figure 7. Distribution map of *Bufo bufo* in the studied area.




Figure 8. Distribution map of *Bufotes viridis* in the studied area.

of tree frogs spawned in someone's garden in a water barrel used for watering the crops. Although some tadpoles were lost as a result of the water being taken out for irrigation, a fair number of them reached maturity. Because of the nocturnal lifestyle of this amphibian and its cryptic colours, many of the identifications were made by sound. The telling apart of this species from *Hyla arborea* Linnaeus, 1758 was based on the distribution according to Stöck et al. 2012. (Figure 9)

Family Ranidae
Pelophylax sp.

Because of the difficulty of identifying these species without capturing them (Cogălniceanu 2002) and their capture is many times impossible, I treated them as the *Pelophylax* complex. Despite that, out of the two species and one klepton of water frogs in Romania, I managed to identify with certainty the marsh frog (*Pelophylax ridibundus* (Pallas, 1771)). However, it is plausible that there might also be other species present. Further research is needed. I should note that the University of Pitești has in its collection 2 water frogs labeled as "*Rana esculenta*" collected in 1965 in Pitești by Gh. Stănescu, but I have not identified the specimens personally. The water frogs are very common in certain places. They can be found easily in ponds and streams at the edge of the Trivale Forest, but also on the course of the Argeș River and in temporary ponds on its banks. (Figure 10)




Figure 9. Distribution map of *Hyla orientalis* in the studied area.




Figure 10. Distribution map of *Pelophylax* in the studied area.

Rana dalmatina Fitzinger in Bonaparte, 1839

The agile frog is an abundant species that was found in many parts of the Trivale Forest. Being a typically woodland animal, it is to be expected to find it mostly in forests and near them, but I have one rather peculiar record of a roadkill individual on the bank of the Argeș River on March 30th 2019, the closest thing to a forest in the area being the Lunca Argeșului Park, which is a pretty well-kept park with planted trees of various species. (Figure 11)

Class Reptilia

Order Testudines

Family Emydidae

Emys orbicularis (Linnaeus, 1758)

The European pond terrapin was found on the Argeș River, including the Prundu Lake and in the Ștrand Park, a city park built on the bank of the Argeș River. One somewhat curious sighting was a juvenile terrapin found in a small pond at the edge of the Trivale Forest, far away from the river. Its presence there could have been caused by a release by some local, but it is also very likely to be a natural occurrence. Inside the city I found a lot of specimens of this species translocated by humans in an




Figure 11. Distribution map of *Rana dalmatina* in the studied area.

artificial pond in a park called Expo Park (located 44.862600, 24.870900). Together with Elena G., a volunteer at the NGO “Wilderness Research and Conservation” we obtained permission from the park’s administration to go and capture the native species in order to release them into the wild. On July 31st 2021 there were rescued 36 European pond terrapins. Because of the conditions in the park, some of the terrapins had health problems, such as eye problems due to the water condition, malnutrition (lack of food, their diet was formed in part out of the junk food thrown by visitors in water) (Rangel-Mendoza et al. 2014) and wounds on the legs and plastron, possibly caused by the rough materials (concrete) that were the only available surfaces to bask on. After the period of rehabilitation, they were released into natural habitats. Due to the anthropic origin of those specimens and their subsequent release into a natural waterway, I decided not to include the location on the map. (Figure 12)

Trachemys scripta (Thunbegr in Schoepff, 1792)

The pond slider is an American species of turtle that used to be very popular in the pet trade. It originates from Southwestern USA and for many years it was sold in many petstores for fairly small prices. Unfortunately, people buy these cheap animals and, as they grow, realize they cannot take care of them anymore, resulting in their abandonment. It is listed as one of the species on the European list of invasive species




Figure 12. Distribution map of *Emys orbicularis* in the studied area.

(<https://www.invazive.ccmesi.ro/>). Many exotic turtles, especially pond sliders, were dumped in the artificial park mentioned previously, but some have been released right in the Argeș River and in the Ștrand Park, from where more Sliders can find their way into the natural water flow. I have observed two subspecies of pond sliders: the red-eared slider (*Trachemys scripta elegans* Wied-Neuwied 1838) and the yellow-bellied slider (*Trachemys scripta scripta* Thunberg in Schoepff 1792). In the habitats where both *Trachemys scripta* and *Emys orbicularis* are present, the exotic species may compete with the native species for the basking spots (Polo-Cavia et al. 2015). Being a bigger species that can also carry dangerous disease for our native turtles, they have the potential to cause harm (Cadi and Joly 2004; Meyer et al. 2015; Iglesias et al. 2015). Even more so as the river grants them access to disperse into more natural habitats. (Figure 13)

Order Squamata
Family Lacertidae
Lacerta agilis Linnaeus, 1758

I only observed the sand lizard in one area, and that is the Ștrand Park, located on the bank of the Argeș River. A somewhat scarce lizard, compared to the other species




Figure 13. Distribution map of *Trachemys scripta* in the studied area.

that I identified, it was found predominantly in the bushy part of the park, sharing the habitat with wall lizards and eastern green lizards. (Figure 14)

Lacerta viridis (Laurenti, 1768)

The eastern green lizard was more frequently observed than its relative, the sand lizard. I identified this species in numerous spots such as inside the Trivale Forest and at its edges, the banks of the Argeş River, as well as inside the city, around the vegetation between apartment blocks. (Figure 15)

Podarcis muralis (Laurenti, 1768)

Without a doubt, the common wall lizard was the most numerous species of reptile that I managed to observe. On October 7th 2018 I was able to count around 94 individuals in one outing, in the course of 1h30min. It was identified in a few places that can, more or less, be connected by following the railway. There are a few spots far from the said railway, but tracing back, they all seem to be “sprouting” from there. This is why I theorize that the population of wall lizards in Piteşti has expanded all the way using trains and the railway (Hedeen and Hedeen 1999; Covaciu-Marcov et al. 2006; Gherghel et al. 2009; Gherghel and Tedrow 2019). The man-made




Figure 14. Distribution map of *Lacerta agilis* in the studied area.




Figure 15. Distribution map of *Lacerta viridis* in the studied area.

habitats are perfect for this species: it was very numerous in places like abandoned train wagons, previously mentioned railways, concrete walls and so on. It must be mentioned that there were observed some individuals that resembled very much the Italian subspecies: *Podarcis muralis nigriventris* (Bonaparte, 1838) but their capture was impossible and so we cannot talk about a 100% accurate identification. If indeed we are talking about the *nigriventris* subspecies, this could be a result of accidental introduction (Michaelides et al. 2015). More research is needed on this aspect of alien herpetofauna (Damas-Moreira et al. 2020). (Figure 16)

Family Anguidae
Anguis colchica (Nordmann, 1840)

The eastern slow worm was only found in two areas of the Trivale Forest. It is of no surprise, as this is a woodland animal. One of them might have been predated and another three were found as roadkill, as they were recorded either on the road or on a path in the forest frequented by joggers and dog walkers. Only one living individual was observed. (Figure 17)




Figure 16. Distribution map of *Podarcis muralis* in the studied area.




Figure 17. Distribution map of *Anguis colchica* in the studied area.

Family Colubridae
Coronella austriaca Laurenti, 1768

The smooth snake has proven fairly cryptic. It was identified in several places, mostly in and around the Trivale Forest, but there is also a record sent in by a local who filmed one in their garden, far from the rest of the records, but still somewhat close to a small tree-covered area. (Figure 18)

Family Natricidae
Natrix natrix (Linnaeus, 1758)

The grass snake is the most common species of snake in the area the study was conducted. Being a reptile that dwells in wet places and feeds mostly on amphibians, it should be of no surprise that it was observed on the banks of the Argeș River, in parks with ponds and channels, around temporary water bodies with *Bombina variegata* in the forest and small ponds in rural areas. (Figure 19)




Figure 18. Distribution map of *Coronella austriaca* in the studied area.




Figure 19. Distribution map of *Natrix natrix* in the studied area.

Natrix tessellata (Laurenti, 1768)

The records of dice snakes in Argeș County are very rare (Conete D., pers. communication), this presented here being, to my knowledge, the first published record in the county. I was not the finder, instead I got this observation from a post on social media made by a local. The spot where this snake was found is inside the Ștrand Park in late October 2019. Its habitat is most likely the cemented banks of the Argeș River. The follow-up of the Dice Snake that was photographed by the aforementioned local is unknown. Nonetheless, it is an important information regarding the distribution of the species in the county and, to my knowledge, this represents the first published record of *Natrix tessellata* in Argeș county. (Figure 20)

Family Viperidae *Vipera berus* (Linnaeus, 1758)

There have been unverified reports of adders - *Vipera berus* (Linnaeus, 1758) by the locals, as well as possible snakebite cases in dogs. This, together with the fact that the species was confirmed in locations surrounding Pitești (just a few kilometers from the studied area) and some of the habitats visited matches pretty well the habitat of this species, makes me believe that the existence of a population is probable, even




Figure 20. Distribution map of *Natrix tessellata* in the studied area.

though no individual was observed during the survey. For this reason, the species is not included in the list of identified species of herpetofauna.

Conservation

The expansion of the urban and suburban areas. One of the most important threat to wild reptiles and amphibians is habitat loss (Brewster et al. 2018). In the last years a lot of new houses have been built in the meadows surrounding the woodland, and parts of the forest have been cleared for the same purpose. This puts at risk species such as *Triturus cristatus*, *Lissotriton vulgaris*, *Salamandra salamandra*, *Bombina variegata*, *Coronella austriaca*, *Natrix natrix* and *Anguis colchica* mainly due to the habitat loss (Brewster et al. 2018).

Illegal collecting for the pet trade. Many species are put at risk because of illegal collecting and faulty exploitation (Altherr and Lameter 2020). The most popular species of herpetofauna that is being captured illegally from the wild is *Emys orbicularis* (Mărginean et al. 2018), but there was one instance of people being observed capturing and taking home *Podarcis muralis* individuals. The *Emys*

orbicularis found in the Expo Park are a prime example of their popularity when it comes to being taken home as pets by people.

Heavy traffic. Each year, a significant number of reptiles and amphibians fall victim to cars on the roads going through their habitat (Fahrig et al. 1995; Colino-Rabanal and Lizana 2012), especially during the breeding migration of amphibians. Some species that have been observed as roadkill more numerous are *Bufo viridis*, *Bufo bufo*, *Anguis colchica* and *Natrix natrix* (although the latter might have also included instances of deliberate killing).

Deliberate killing by locals. Many reptiles and amphibians are killed by locals due to myths, misbeliefs, fear or ignorance (Fuhn 1969, Marshall et al. 2020). The most common victims are species of snakes, but also *Anguis colchica* due to the confusion of this species with a snake (Fuhn 1969) and even *Lacerta viridis* (pers. observation). There is a long road to go until people finally understand the importance of these animals, but it is possible with education.

Invasive species. The impact of a newly introduced species in an ecosystem on the native species can be huge, at times leading to extinction (Clavero and Garcia-Berthou 2005). Some examples of alien species that have the potential to have an impact on the local herpetofauna are *Trachemys scripta* (which's effect has been discussed earlier) and *Felis catus* (Woods et al. 2003, Loss et al. 2013) which has been observed preying on *Rana dalmatina*, *Lacerta viridis* and *Natrix natrix* (pers. observation). Measures are advised to be taken in order to limit the spreading of this invasive species, such as fining more harshly the abandonment of pets, supporting and facilitating the capturing of invasive species and setting up centers to take sliders into their custody.

Out of all the 18 species recorded in the study area, 13 are included in The Red Book of Romanian Vertebrates, 17 in the Berne Convention on the Conservation of European Wildlife and Natural Habitats and in 16 in Order 57/2007 issued by the Government of Romania and one on the list of invasive alien species of European Union concern (Table 4).

It should be noted that at the time the Convention took place and the law and the book were written, *Hyla orientalis* was part of *Hyla arborea* (Stöck et al. 2012), *Pelophylax ridibundus* was *Rana ridibunda* (Dubois and Ohler 1994), *Bufo viridis* was *Bufo viridis* (Dufrenoy et al. 2019), *Lissotriton vulgaris* was named *Triturus vulgaris* (Titus and Larson 1995) and *Anguis colchica* belonged to *Anguis fragilis* (Gvozdík et al. 2010). I used the updated taxonomy based on the new genetic-based studies in this paper and placed the protection status accordingly.

Table 4. Status of protection of the species of amphibians and reptiles recorded

Species	Berne Convention on the Conservation of European Wildlife and Natural Habitats	Order 57/2007	The Red Book of Romanian Vertebrates	The IUCN Red List of Threatened Species
<i>Salamandrina salamandrina</i>	Berne III	Order 57/2007 annex 4B	VU	LC
<i>Lissotriton vulgaris</i>	Berne III	Order 57/2007 annex 4B	NT	LC
<i>Triturus cristatus</i>	Berne II	Order 57/2007 annex 3 & 4A	VU	LC
<i>Bombina variegata</i>	Berne II	Order 57/2007 annex 3 & 4A	NT	LC
<i>Bufo bufo</i>	Berne III	Order 57/2007 annex 4B	NT	LC
<i>Bufo viridis</i>	Berne II	Order 57/2007 annex 4A	NT	LC
<i>Hyla orientalis</i>	Berne II	Order 57/2007 annex 4A	VU	LC
<i>Pelophylax ridibundus</i>	Berne III	Order 57/2007 annex 5A		LC
<i>Rana dalmatina</i>	Berne II	Order 57/2007 annex 4A	VU	LC
<i>Enys orbicularis</i>	Berne II	Order 57/2007 annex 3 & 4A	VU	NT
<i>Trachemys scripta</i> *				LC
<i>Lacerta agilis</i>	Berne II	Order 57/2007 annex 4A		LC
<i>Lacerta viridis</i>	Berne II	Order 57/2007 annex 4A		LC
<i>Podarcis muralis</i>	Berne II	Order 57/2007 annex 4B	VU	LC
<i>Anguis colchica</i>	Berne III	Order 57/2007 annex 3B	VU	LC
<i>Coronella austriaca</i>	Berne II	Order 57/2007 annex 4A	VU	LC
<i>Natrix natrix</i>	Berne III			LC
<i>Natrix tessellata</i>	Berne II	Order 57/2007 annex 4A	NT	LC

*List of invasive alien species of European Union concern

Conclusions

In the city of Pitești and the area immediately neighboring I have recorded nine species of amphibians, belonging to five families and two orders, making up 47.36% of the species in the Romanian amphibian fauna (Fuhn 1960; Stöck et al. 2012), as well as nine species of reptiles belonging to five families, three clades and two orders, the eight native reptiles making up 33.33% of the species in the Romanian reptile fauna (Fuhn and Vancea 1961). Apart from better defining the known range of already acknowledged species of the area, the study also brought more light to the expansion of allochthonous reptiles like *Trachemys scripta*. Overall, the paper hereby present fills a few gaps in our knowledge about the fauna of Argeș County, even if a few more situations remain to be researched further.

Acknowledgements

I am indebted to the curators of the Argeș County Museum and the professors from the University of Pitești for giving me acces to their collections and data, as well as to Denisa Conete, Iulian Gherghel, Raluca Melenciuc, Tiberiu Sahlean, Alexandru Strugariu, for advice on writing this paper, to Ștefan Bădoi, Dinu Berevoianu, Georgian Neacșu, Anca Pavel, Tiberiu Sahlean, Raluca Udrea, for assistance in fieldwork, to Tiberiu Sahlean for providing the distribution maps modeled in ArcGIS, and to Iulian Gherghel, Raluca Melenciuc, Alexandru Pintilioaie, Tiberiu Sahlean and Alexandru Strugariu for reviewing this manuscript.

References

- Altherr S, Lameter K (2020) The Rush for the Rare: Reptiles and Amphibians in the European Pet Trade. *Animals* 10: 2085. doi:10.3390/ani10112085.
- Bickford D, Howard SD, Ng DJ, Sheridan JA (2010) Impacts of climate change on the amphibians and reptiles of Southeast Asia. *Biodiversity and Conservation* 19: 1043–1062 DOI 10.1007/s10531-010-9782-4.
- Brewster CL, Beaupre SJ, Willson JD (2018) Habitat Loss and Local Extinction: Linking Population Declines of Eastern Collared Lizards (*Crotaphytus collaris*) to Habitat Degradation in Ozark Glades. *Journal of Herpetology* 52(3): 352–360.
- Cadi A, Joly P (2004) Impact of the introduction of the red-eared slider (*Trachemys scripta elegans*) on survival rates of European pond turtle (*Emys orbicularis*). *Biodiversity and Conservation* 13: 2511–2518.
- Clavero M, García-Berthou E (2005) Invasive species are a leading cause of animal extinctions. Institute of Aquatic Ecology, University of Girona, E-17071 Girona, Spain.

- Cogălniceanu D (2002) Amfibienii din România. Ghid de teren. Naturalia Practica no. 5. Colecția de Biologie-Ecologie, Universitatea din București. Editura Ars Docendi, 1–41. [in Romanian]
- Cogălniceanu D, Székely P, Samoilă C, Iosif R, Tudor M, Plăiașu R, Stănescu F, Rozyłowicz L (2013) Diversity and distribution of amphibians in Romania. *ZooKeys* 296: 35–57. 10.3897/zookeys.296.4872.
- Colino-Rabanal VJ, Lizana M (2012) Herpetofauna and roads: a review. *Basic and Applied Herpetology* 26: 5–31.
- Covaciu-Marcov SD, Bogdan HV, Ferenti S (2006) Notes regarding the presence of some *Podarcis muralis* (Laurenti 1768) populations on the railroads of western Romania. *North-Western Journal of Zoology* 2(2): 126–130
- Covaciu-Marcov SD, Sas I, Ilieș A (2010) *Pelophylax lessonae* (Amphibia) in Râul Doamnei, Argeș County, Romania. How have we arrived here? *Bihorean Biologist* 4(1): 83–87.
- Covaciu-Marcov SD, Cicort-Lucaci AȘ, Telcean IC, Pal A, Sas-Kovács I (2014) Some notes on the herpetofauna from Vâlsan River Natural Protected Area, Romania. *Carpathian Journal of Earth and Environmental Sciences* 9(3): 171–176.
- Damas-Moreira I, Riley JL, Carretero MA, Harris DJ, Whiting MJ (2020) Getting ahead: exploitative competition by an invasive lizard. *Behavioral Ecology and Sociobiology* 74: 117.
- Delaney K, Busteed G, Fisher R, Riley S (2021) Reptile and Amphibian Diversity and Abundance in an Urban Landscape: Impacts of Fragmentation and the Conservation Value of Small Patches. *Ichthyology & Herpetology* 109. 10.1643/h2019261.
- Dubois A, Ohler A (1994) Frogs of the subgenus *Pelophylax* (Amphibia, Anura, genus *Rana*): A catalogue of available and valid scientific names, with comments on name-bearing types, complete synonymies, proposed common names, and maps showing all type localities. *Zoologica Poloniae* 39(3): 139–204.
- Dufresnes C, Mazepa G, Jablonski D, Caliani Oliveira R, Wenseleers T, Шабанов Д, Auer M, Ernst R, Ramírez-Chaves H, Mulder K, Simonov E, Tiutenko A, Kryvokhyzha D, Wennekes P, Zinenko O, Korshunov O, Al-Johany A, Peregontsev E, Masroor R, Litvinchuk S (2019) Fifteen shades of green: The evolution of *Bufo* toads revisited. *Molecular Phylogenetics and Evolution* 141. 10.1016/j.ympev.2019.106615.
- Edgar P, Foster J, Baker J (2010). *Reptile Habitat Management Handbook*. Amphibian and Reptile Conservation, Bournemouth.
- Fahrig L, Pedlar JH, Pope SE, Taylor PD, Wegner JF (1995) Effect of road traffic on amphibian density. *Biological Conservation* 73: 177–182.
- Farinha-Marques P, Lameiras JM, Fernandes C, Silva S, Guilherme F (2011) Urban biodiversity: a review of current concepts and contributions to multidisciplinary approaches. *Innovation: The European Journal of Social Science Research* 24(3): 247–271. doi:10.1080/13511610.2011.592062.
- Fuhn IE (1960) Fauna Republicii Populare Române. Vol. XIV. Fascicula 1: Amphibia. București, Romania. Editura Academiei Republicii Populare Române, 288. [in Romanian]

- Fuhn IE, Vancea Ș (1961) Fauna Republicii Populare Române. Vol. XIV. Fascicula 2: Reptilia (Testoase, șopârle, șerpi). București, Romania. Editura Academiei Republicii Populare Române, 352. [in Romanian]
- Fuhn IE (1969) Broaște, șerpi, șopârle. București, Romania. Editura Științifică, 244 pp. [in Romanian]
- Gherghel I, Strugariu A, Sahlean TC, Zamfirescu O (2009) Anthropogenic impact or anthropogenic accommodation? Distribution range expansion of the common wall lizard (*Podarcis muralis*) by means of artificial habitats in the north-eastern limits of its distribution range. *Acta Herpetologica* 4(2): 183–189.
- Gherghel I, Tedrow R (2019) Manmade structures are used by an invasive species to colonize new territory across a fragmented landscape. *Acta Oecologica* 101. 10.1016/j.actao.2019.103479.
- Gibbons JW, Scott DE, Ryan TJ, Buhlmann KA, Tuberville TD, Metts BS, Greene JL, Mills T, Leiden Y, Poppy S, Winne CT (2000) The Global Decline of Reptiles, Déjà Vu Amphibians: Reptile species are declining on a global scale. Six significant threats to reptile populations are habitat loss and degradation, introduced invasive species, environmental pollution, disease, unsustainable use, and global climate change. *BioScience* 50(8): 653–666.
- Guoždík V, Jandzik D, Lymberakis P, Jablonski D, Moravec J (2010) Slow worm, *Anguis fragilis* (Reptilia: Anguidae) as a species complex: Genetic structure reveals deep divergences. *Molecular Phylogenetics and Evolution* 55: 460–472.
- Hamer AJ, McDonnell MJ (2010) The response of herpetofauna to urbanization: Inferring patterns of persistence from wildlife databases. *Austral Ecology* 35: 568–580.
- Hedeen SE, Hedeen DL (1999) Railway-aided dispersal of an introduced *Podarcis muralis* population. *Herpetological Review* 30(1).
- Iftime A, Iftime O (2014) Notes on the herpetofauna of the Leaota Mountains, a “wildlife corridor” area. *North-Western Journal of Zoology* 10(1): S33–S37.
- Iglesias R, García-Estévez JM, Ayres C, Acuña A, Cordero-Rivera A (2015) First reported outbreak of severe spirorchidiasis in *Emys orbicularis*, probably resulting from a parasite spillover event. *Diseases of Aquatic Organisms* 113(1):75–80.
- Loss SR, Will T, Marra PP (2013) The impact of free-ranging domestic cats on wildlife of the United States. *Nature Communications* 4: 1396.
- Marshall BM, Strine C, Hughes AC (2020) Thousands of reptile species threatened by under-regulated global trade. *Nature Communications* 11: 4738. <https://doi.org/10.1038/s41467-020-18523-4>.
- Mayani-Parás F, Botello F, Castañeda S, Sánchez-Cordero V (2019) Impact of Habitat Loss and Mining on the Distribution of Endemic Species of Amphibians and Reptiles in Mexico. *Diversity* 11: 210. <https://doi.org/10.3390/d11110210>.
- Mărginean GI, Gherman E, Sos T (2018) The illegal internet based trade in European pond turtle *Emys orbicularis* (Linnaeus, 1758) in Romania: a threat factor for conservation. *North-Western Journal of Zoology* 14(1): 64–70.
- Meyer L, Preez L, Bonneau E, Héritier L, Franch M, Valdeón A, Sadaoui A, Kechemir-Issad N, Palacios C, Verneau O (2015) Parasite host-switching from the invasive American

- red-eared slider, *Trachemys scripta elegans*, to the native Mediterranean pond turtle, *Mauremys leprosa*, in natural environments. *Aquatic Invasions* 10(1): 79–91.
- Michaelides SN, While GM, Zajac N, Uller T (2015) Widespread primary, but geographically restricted secondary, human introductions of wall lizards, *Podarcis muralis*. *Molecular Ecology* 24: 2702–2714.
- Polo-Cavia N, Lopez P, Martin J (2015). Interference competition between native Iberian turtles and the exotic *Trachemys scripta*. *Basic and Applied Herpetology* 28. 10.11160/bah.13014.
- Rangel-Mendoza J, Sánchez-González IA, López-Luna M, Weber M (2014) Health and Aquatic Environment Assessment of Captive Central American River Turtles, *Dermatemys mawii*, at Two Farms in Tabasco, Mexico. *Chelonian Conservation and Biology* 13: 96–106. 10.2744/CCB-1040.1.
- Speybroeck J, Beukema W, Bok B, Van Der Voort J (2016) Field Guide to the Amphibians and Reptiles of Britain and Europe. London UK. Bloomsbury Natural History, 432.
- Stöck M, Dufresnes C, Litvinchuk SN, Lymberakis P, Biollay S, Berroneau M, Borzée A, Ghali K, Ogielska M, Perrin N (2012) Cryptic diversity among Western Palearctic tree frogs: Postglacial range expansion, range limits, and secondary contacts of three European tree frog lineages (*Hyla arborea* group). *Molecular Phylogenetics and Evolution* 65(1): 1–9.
- Strugariu A, Zamfirescu ȘR, Gherghel I (2009) First record of the adder (*Vipera berus berus*) in Argeș County (Southern Romania). *Bihorean Biologist* 3(2): 163–166.
- Stuart SN, Chanson JS, Cox NA, Young BE, Rodrigues AS, Fischman DL, Waller RW (2004) Status and trends of amphibian declines and extinctions worldwide. *Science* 306(5702):1783–6. doi: 10.1126/science.1103538.
- Titus TA, Larson A (1995) A molecular phylogenetic perspective on the evolutionary radiation of the salamander family Salamandridae. *Systematic Biology* 44: 125–151.
- Török Z, Ghira I, Sas I, Zamfirescu Ș (2013) Ghid sintetic de monitorizare a speciilor comunitare de reptile și amfibieni din România. Tulcea, Romania. Editura Centrul de Informare Tehnologică “Delta Dunării”, 116. ISBN 978-973-88117-6-8. DOI: 10.7427/DDI.B.01.2013. [in Romanian]
- Woods M, McDonald RA, Harris S (2003) Predation of wildlife by domestic cats *Felis catus* in Great Britain. *Mammal Review* 33(33): 174–188.